



VER THE PAST SEVERAL YEARS budgets for turfgrass maintenance have been reduced dramatically. Athletic field managers have all had to adjust to the struggles associated with maintaining high quality turf with less money. One could argue that sports turf managers are affected more than any other sector of the industry during these trying times. While dollars for field maintenance continue to decline, demands on field use do not. To conquer these challenges from a weed control perspective, sports turf managers must comprehensively evaluate their management programs to determine where valuable budget dollars should be spent. The steps presented below can be used as a guide for making weed control decisions on a restricted budget.

PRIORITIZE

The first step in effectively controlling weeds on a restricted budget is to realize that any program implemented most likely will not be comprehensive. While the entire facility may have been treated in the past, with fewer dollars in the budget it is critically important that sports turf managers determine what parts of their facility need to look the best (i.e., be free of weeds). Many sports turf managers grade these areas. For example, game fields for varsity sports may be deemed A-level areas, practice fields may be deemed B-level areas, and those used for band or physical education class may be deemed C-level areas.

TIMING

Once key areas have been identified, it is important to determine when these areas need to look the best. If fall sports predominate, weed control in early spring becomes less important. This especially rings true for those managing bermudagrass fields in the transition zone. There are significant costs associated with fall overseeding including the cost of seed, fertilizer, water (in some cases), labor for mowing, and herbicides for weed control. Chemically removing the overseeded turf in spring can cost as much as \$150 per acre, in addition to the cost of selective winter annual broadleaf and grassy weed control.

On fields with minimal spring play, consider not overseeding in fall and controlling weeds with a non-selective herbicide in spring while the bermudagrass is still dormant. This strategy will provide economical control of winter annual broadleaf and grassy weeds with a single herbicide application. Lack of competition from the overseeded turf will also allow bermudagrass to recuperate from fall traffic more efficiently in spring. Research conducted at the University of Tennessee in 2009 found programs delivering non-selective herbicide applications to dormant bermudagrass to provide more effective and economical weed control than those incorporating fall overseeding and selective herbicide applications.

Proper herbicide selection depends on many factors including the weed requiring control, the stage of weed maturity, and the species of turfgrass in which the weed has invaded.

PRESSURE

Another important step in controlling weeds with less money is to determine the amount of weed pressure that warrants an herbicide application. This is directly related to the expectations of field users and the scope of the budget provided. While a 100% weed-free stand may have been the standard in the past, budgets may require that this be shifted to 85% or 90%. Another option is to ensure that A-level areas are 100% weedfree, while B and C-level areas remain at 70-80%. These adjustments may help stretch dollars for weed control further.

CULTURAL PRACTICES

A dense stand of vigorously growing turf is the best defense against weed invasion. Weeds only invade turf stands if there is a void in the canopy in which to do so. With fewer budgetary dollars available for weed control, sports turf managers should thoroughly evaluate their cultural practices to ensure that everything possible is being done to maintain turf density under traffic.

Practices such as selecting traffic tolerant cultivars, applying fertilizers to meet soil test recommendations, sufficiently irrigating to meet plant needs, mowing at a proper height of cut, aerfiying regularly, and applying plant growth regulators all serve to maximize turf density and consequently reduce the likelihood of weed infestations. In a world of smaller budgets, implementing cultural practices to maximize turfgrass density should be thought of as preventative weed control.

CALIBRATION

Calibrating spraying equipment is probably the easiest way to save money when budget dollars are limited. Studies have shown that over 80% of licensed pesticide applicators have calibration and/or mixing errors greater than 5%. These errors have been found to range from a 40% under application to a 60% over application, resulting in over 4 million dollars in lost revenue.

Sprayers should be calibrated at least one time per growing season, with nozzles changed regularly. Taking the time to check that spraying equipment is calibrated and operating properly will ensure that dollars are not wasted on improper herbicide applications.

SELECTING THE CORRECT PRODUCT

Proper herbicide selection depends on many factors including the weed requiring control, the stage of weed maturity, and the species of turfgrass in which the weed has invaded. New herbicides are entering the marketplace with a focus on broadening the weed control spectrum of a single herbicide application. These products typically combine two or more herbicides that have been sold individually into a single formulated product. Examples include Solitare (sulfentrazone + quinclorac) and Onetime (quinclorac + MCPP + dicamba). Both of these products offer postemergence control of crabgrass and various broadleaf weeds with a single application. Using a product with a wide weed control spectrum will reduce the number of herbicide applications required during the season.

PREPARING FOR LOSS OF MSMA

The loss of an effective, economical postemergence herbicide like MSMA will make weed control even more difficult with restricted budgets. Questions about legal applications of MSMA still linger throughout the industry. According to the EPA, registrants of MSMA for use on sports turf lost the ability to sell the product on December 31, 2009; however, distributors will be able to sell products purchased before December 31, 2009 until December 31, 2010. After December 31, 2010 existing stocks of MSMA can legally be used for weed control on sports fields until they are exhausted provided that these uses comply with the EPAapproved label and labeling of the affected product.

Considering that the window to legally apply MSMA is closing by the minute, sports field managers challenged with dallisgrass infestations should place top priority on controlling these problems as soon as possible. Researchers at the University of Tennessee are continually evaluating new compounds, as well as combinations of existing compounds, that will help soften the blow of the EPA restrictions on MSMA. To follow the latest research being conducted at the University of Tennessee, visit http://tennesseeturfgrassweeds.org.

All aspects of turfgrass management, including weed control, become challenging following budget cuts. Taking the time to evaluate why certain practices have been implemented in the past should help identify the key components of the maintenance program requiring attention (and budget dollars) in the future.

Dr. James T. Brosnan is assistant professorturfgrass weed science at the University of Tennessee; Greg Breeden is weed science extension assistant at UT.



Dr. James Brosnan, University of Tennessee, Technical Editor of *SportsTurf*.



SOIL TESTING IS NOT ABOUT FERTILITY At least not exclusively!

OR OVER TWENTY YEARS I have been staring at soil tests, writing reports and recommendations. And for over twenty years I have seen, sports turf managers, and golf course superintendents make changes to their properties. Clearly they have produced better turf but more importantly they have made changes to their soil profile. They have improved drainage, strengthened rooting, increased plant recovery and they reduced fertility inputs. Why? Because they have soil tested and their soil testing programs have helped them to discover that improving the physical structure and biological profile of the soil is the precursor to good soil and plant fertility.

The four basics of good soil management are: Air Management, Water Management, Digestion Management and finally Nutrient Management, in that order. Air space in the soil allows for good water movement, without air in the soil roots will not survive and beneficial soil micro-organisms will not proliferate. With too much water an anaerobic condition will prevail and both roots and microbes will die off. With good air and water movement through the soil profile you will create the environment where beneficial soil microbes can freely multiply and in the process carbon is digested down to humus — the ultimate break down of organic matter in the soil. This humus is then used as a microbial food source to generate even more biological activity. It is the back bone of the carbon to nitrogen cycle in the soil, producing free and needed nitrogen to the plant without all the side effects of synthetic applications. If and only if the environment for microbial digestion is present in the soil will nutrient mobility take place. Microbes "eat at the table first" and all nutrient mobility from the soil is done through microbial degradation. A good soil testing protocol allows the turf manager to manipulate the soil physically, by balancing basic cations and relaxing soil particles just enough to allow better air and water movement through the soil profile. This builds a better environment for soil micro-organisms to do their job of digesting carbon, releasing nutrients into the root zone and providing buffers for water, temperature and pathogen attack. So, if 75% of good soil management deals with air, water and carbon digestion (physics and biology), is soil testing really about fertility?

The intention is to change soil physics and improve the environment for soil biology. After more than twenty years of soil testing, tens of thousands of soil tests on hundreds of sites, I have never seen the process fail if the turf manager sticks to the basics. I have, however, repeatedly been told that it cannot work, "you can't change soils physically by changing soils chemically", "there is no research to prove this", "an acre inch is two million pounds of soil much too much to change..." The truth is you can change soils physically by manipulating the soil chemically and you better if your soils are out of balance and you hope to produce better turf with fewer inputs. We are not changing clay to sand or sand to silt and we are not concerned about the entire two million pounds of soil in the acre inch, we are only concerned about the four thousand pounds or so that make up the nutrient profile particularly calcium, magnesium and potassium that affect soil flocculation which impacts soil biology and that we can change. What we are really focused on is Biological Soil Management.

According to the working models presented by Dr.

After more than twenty years of soil testing, tens of thousands of soil tests on hundreds of sites, I have never seen the process fail if the turf manager sticks to the basics.



Albrecht at the University of Missouri, and Dr. Bear at Rutgers University we are looking to balance the basic soil cations and striving for a profile that presents roughly 68% calcium, 12% magnesium, 5% potassium, 2 % sodium, 3% trace elements and 10% hydrogen. When a colloidal soil moves to these percentages the soil physically works better. Air and water move through the profile more efficiently building an environment where soil biology explodes and the nitrogen cycle increases. In this model if we can achieve 10% hydrogen we will always have a pH of 6.3 which is the point at which we find maximum potential nutrient mobility across the entire nutrient board. These percentages are found in the base saturation section of a complete base saturation soil test and can easily be manipulated by adding what is deficient and knocking off what is excessive. Unfortunately, not all soil tests show a complete base saturation where all six cation groups are present, calcium, magnesium, potassium, trace elements, sodium and hydrogen. Too often a soil testing laboratory will show a limited spectrum of soil cations or they will add up to something more that 100% which is not a true base saturation and makes building a quality recommendation all most impossible.

The following base saturation was found on a soil test of a sports turf client: 35% Ca; 45% Mg; 2% K; 5% Na; 3% traces and 10% hydrogen. He was not able to aerify the soil because it was too hard

Job Nar	ame STB Hills CC Date 1/1/2007									
Compa	any Logan Labs 888-494-7645 Submitted By									
Sample	Location		Ideal							
Sample	ID		1	2	3	4	5			
Lab Nui	mber									
Sample Depth in inches			6	6	6	6	6			
Total Exchange Capacity (M. E.)			12.00	2.74	8.89	21.03	12.80			
pH of Soil Sample			6.30	5.60	6.50	7.20	6.30			
Organic Matter, Percent			3-5	0.80	3.54	1.40	1.20			
SNO	SULFUR:	p.p.m.	25-50	8	55	18	12			
ANIO	Mehlich III Phosphorous:	as (P ₂ O ₅) lbs / acre	250-500	232	284	250	242			
SN	CALCIUM: Ibs / acre	Desired Value Value Found Deficit	3264 3264 -0	745 459 -286	2418 2308 -110	5720 4380 -1340	3482 1777 -1705			
ABLE CATIO	MAGNESIUM: Ibs / acre	Desired Value Value Found Deficit	346 348 -0	200 138 -62	256 355	606 2038	369 1369			
EXCHANGE	POTASSIUM: Ibs / acre	Desired Value Value Found Deficit	468 468 -0	163 33 -130	277 279	552 248 -304	396 255 -141			
	SODIUM:	lbs / acre	20-50	30	84	176	153			
%	Calcium (60 to 70%)		68.0	41.88	64.90	53.07	34.71			
NOI	Magnesium (10 to 20%)		12.0	20.99	16.64	40.38	44.55			
IRAT	Potassium (2 to 5%)		5.00	1.54	4.02	1.51	2.55			
SATI	Sodium (.5 to 3%)		3.00	2.38	2.05	1.82	2.60			
ASE	Other Bases (Variable)		2.00	6.20	4.90	4.20	5.10			
8	Exchangable Hydrogen (10 to 15%)		10.00	27.00	7.50	0.00	10.50			
TS	Boron (p.p.m.)		1.20	0.02	0.48	0.7	0.66			
MEN	Iron (p.p.m.)		150	175	319	740	540			
	Manganese (p.p.m.)		40	2	2.09	145	26			
ACE	Zinc (p.p.m.)		10	1.8	7 44	5.5	3.5			
TR	Aluminum (p.p.m.)		<1400	200	419	800	900			
THER										

Logan Labs, LLC

and the metal tines of the aerifier would not penetrate the soil. When it rained the soil would not drain for days, he had very poor rooting, seed germination was weak and he had to apply over 8 pounds of synthetic nitrogen per 1000 square feet to the site each year simply to maintain color. When I asked how his soil was he responded by saying "it's not my soil, I tested it and the pH was 6.3!" He had over a 1700 pound deficiency in calcium, a thousand pound excess of magnesium, low potassium and high sodium. I recommended three applications of high calcium limestone at a rate of one ton per acre, separated by a few months, along with a lot of potassium sulfate. After the second application of lime I called him and asked if he was able to aerify the property and he reported that he could not. So I asked him if he had seen any difference and he did say that he thought they had reformulated his fertilizer because it was working a lot better. With better soil biology fertilizer will work better! One year later he called and told me that he had run his aerifier over this site and did not break a single tine, which was uncommon prior to making the soil adjustments, so he ran the machine off the treated area to an adjacent soil that was not treated and as soon as he hit that soil he broke the machine into pieces! He reduced his nitrogen inputs down to around two pounds per 1000 square feet per year, drainage was no longer a problem, rooting was strong and deep, recovery was excellent and seed germination when he needed it was very good.

Job Na	me STR Hills CC				Da	te 1/1/200	17		
000 Ma		00 101 761E							
Compa	any LOGAIT LADS 6	00-494-7045		IDMITTED BY	1				
Sampi	le Location		Ideal						
Sample ID			1	2	3	4	5		
Lab N	lumber								
Water	- Used		DI	DI	DI	DI	Irrigation		
water Used			6.3	6.5	6.5	7 1	71		
рн			<000	404	450	104	444		
Soluble Salts ppm			<900	134	150	121	114		
Chlori	ide (Cl)	ppm	<50	8	4	8	10		
Bicarb	oonate (HCO3)	ppm	<50	40	176	73	366		
ANIONS	SULFUR	ppm	5-10	10.72	36.67	11.71	9.59		
	PHOSPHORUS	ppm	1-3	0.61	< 0.1	0.38	0.46		
	CALCIUM	ppm	40-60	42.59	25.54	14.34	13.32		
SOLUBLE CATIONS		meq/l	0	2.13	1.28	0.72	0.67		
	MAGNESIUM	ppm	8-12	2.06	5.74	2.11	1.95		
		meq/l	0	0.17	0.48	0.18	0.16		
	POTASSIUM:	ppm	15-20	25.52	10.86	24.51	13.06		
		meq/l	0	0.66	0.28	0.64	0.34		
	SODIUM	ppm	<20	4.56	17.39	5.17	24.82		
		meq/l	0	0.20	0.76	0.22	1.08		
PERCENT	Calcium		55-60	67.34	45.71	40.87	29.64		
	Magnesium		18-20	5.43	17.12	10.02	7.23		
	Potassium		9-12	20.96	10.10	36.29	15.10		
	Sodium		2-8	6.27	27.07	12.81	48.03		
TRACE ELEMENTS	Boron (p.p.m.)		0.1	0.02	0.03	< 0.02	< 0.02		
	Iron (p.p.m.)		0.3	1.35	3.12	1.25	2.55		
	Manganese (p.p.m.)		0.1	0.03	0.15	0.03	< 0.02		
	Copper (p.p.m.)		0.08	0.07	< 0.02	0.08	< 0.02		
	Zinc (p.p.m.)		0.08	0.08	< 0.02	0.11	0.06		
	Aluminum (p.p.m.)		∥ – – – – –	1.86	1.21	1.47	2.94		

Logan Labs, LLC

When staring at as many soil tests as I do it sometimes becomes hard to see the forest through the trees but a client once told me that when the light bulb finally went off for him was when he realized that base saturation was always 100%, no more and no less. If a soil test shows more or less than 100% base saturation it is not a true base saturation test which is the key here. Understanding that it is always 100% makes it easy to see that if something is high it can be exchanged (cation exchange capacity) by replacing it with what is low. In the previous example magnesium was driving the soil and calcium was low so by adding the proper amount and type of calcium we were able to drive off the magnesium which was tightening the soil and affecting soil biology.

Base saturation is only one of many tools that we use to manage soil profiles and on some soils it is not as strong a tool, such as low CEC sand based soils, but the tool can still give us direction. On these sites the sufficiency levels of nutrients is clearly our first focus. I have become a big fan of the water soluble paste extract test to give me even more information and help me answer some questions that may not be easy to answer simply by reviewing the standard soil tests. I use Logan Labs (www.loganlabs.com) for my lab work but there a number of good labs producing complete base saturation soil tests. A complete base saturation soil test will never lie to you it may tell you that sodium is driving the soil like out west or calcium on the calcareous soils of the mid west or even aluminum like in some of the gumbo soils down south. A good soil testing protocol will give you the map and directions on how to affect physics which will affect soil biology. Oh and by the way all of this improves nutrient mobility!

Joel Simmons is the President of EarthWorks Natural Organic Products and Soil First consulting and teaches the Soil First Academy all across North America. He holds a Masters Degree from Penn State University and is a former Penn State County Extension Agent and Instructor of Soils at Rutgers University. He may be reached in front of his computer at joel@soilfirst.com most likely working on yet another client's soil test.



ww.exceneratortun.com [mineralsupport@naisco.cc

"Excellerator provides our fields with exceptional wear tolerance. I would recommend it to any turf manager who wants to improve the peformance and safety of their fields."



Chris Kneale, Bentley College, Waltham, MA

Drought tolerant grass backed by complex technology



Pennington's Extended Root Technology provides increased root depth and mass to create stronger, healthier plants, significantly reducing the amount of water needed to maintain beautiful turf/ It's the perfect solution for your budget and for our environment. Pennington...a trusted brand since 1945.

PENNIÑGTON

www.penningtonseed.com