



Powdery Mildew gets its name from the early stage growth of a gray-white powdery fungus on the leaf surface. In cases of high incidence, entire turf areas may appear dull white. Prevent fungus growth with 1-2 oz. every 14-28 days.



Rust shows up first as light yellow flecks on the grass leaves. If left untreated, leaf surface will eventually rupture and yellow-orange or reddish-brown pustules develop. Preventive treatment calls for 1-2 oz. every 14-28 days.



Spring Dead Spot appears as circular, straw-colored patches on Bermudagrass leaves just as they begin to green-up in spring. The rest of the plant is affected with black to brown dry rot. University research indicates preventive control can be obtained with 4 oz. applied in the fall.*



Brown Patch is recognized by a "frog-eye" pattern formed by the diseased turf encircling green, unaffected plants. For prevention, apply 2-4 oz. every 10-21 days. If disease is already present, apply 2 oz. with a registered contact fungicide.

You treat each disease on its own turf.

inate rate, Banner gives you the precision to treat up to a total of twelve specific turf diseases with their own recommended rates, timing, and frequencies.

Rates that, in many cases, are lower than those that are required with other turf-grass fungicides.

Which means Banner isn't just an intelligent way

to effectively prevent disease, it's an intelligent way to prevent spending more than you have to.

