

MAINTAIN THE DREAM

What Makes Great Teams?

Spirit ~ Strength ~ Endurance ~ Technique
and a **KROMER Maintained Field**

**The B200 – Simply the best
“ALL IN ONE” Athletic Field
Maintainer, ever!**

The only machine that applies
dry lines, wet lines, grooms,
conditions, sprays and will cut
and paint a line at the same time.
Available with 32 attachments.



Traler Groomer™

An economical pull type groomer for use on, baseball, soccer, football, lacrosse, and rugby fields. Conditions, levels, paints lines, and sprays chemicals. Available with multiple attachments for perfect field grooming.



Our Field is Preparing Your Field

Baseball ~ Football ~ Softball ~ Soccer
Rugby ~ Lacrosse ~ Track & Field

*From walk behind liners, to paint mixing stations,
to riders, to groomers – we've got you covered!*

1-800-373-0337
www.kromerco.com

Fill in 121 on reader service form or visit <http://oners.hotims.com/12047-121>

#ST250

LATE FALL N FERTILIZATION IS AN IMPORTANT TIME FOR DEVELOPING A STRONG AND DEEP ROOT SYSTEM



However, the magnitude of leaching was highly dependent on the location. The greatest losses were from the two southeastern coastal sites on Long Island (Riverhead and St. Charles) that had similar soils compared to the more northern Ithaca site, but had half as many days (36 days for Riverhead and St. Charles compared to 70 days for Ithaca) that the average temperature was below freezing. The amount of groundwater recharge during late fall-late to spring period for Riverhead 75-90 % of precipitation (October 15 to May 15) explaining the potential for a large amount of N leaching if water soluble N sources are applied in the late fall. We have observed a lower amount of groundwater recharge in Ithaca.

When N was applied during the growing season, similar results were observed with N source and leaching losses, depending on the year of the study. In years 1 and 2 of the study, where rainfall values were below or close to average amounts, N leaching percent of the amount applied was low and not highly influenced by N source. Year 3, where about 24 % more rainfall than the average, N leaching losses were much greater and soluble N sources like urea and calcium nitrate had large amounts of N leaching. The other N sources had much less leaching ranging from 2 to 7 % of the amount of N that was applied.

In addition to N source and weather, application timing influenced the extent on N leaching losses we observed. The extent of N leaching in the late fall was greatest with water soluble sources and was about four times greater than the N leaching losses observed during the growing season study, even though the growing season study had twice as much N applied over the entire year than the late fall study.

Late fall N fertilization is an important time for developing a strong and deep root system. However, the differences on the extent of freezing conditions in the winter months does play a role in N leaching losses from late fall or winter N applications, especially with water soluble N sources like urea. Thus, areas with less frozen soil conditions in the dormant or slow cool season growth periods when high amounts of rainfall or snow melt is likely to occur, can be prone to excessive N leaching if water soluble N sources like urea, ammonium sulfate, ammonium nitrate and calcium nitrate are used.

Based on the results of this field study, one can conclude that slow release N sources like IBDU, ureaformaldehyde, PCU and SCU and natural organic fertilizers like biosolid appear to significantly reduce the potential of N leaching losses into groundwater compared to the very water soluble calcium nitrate. However, long-term studies are needed to better understand the risk of using slow release N sources as a best management practice to reduce N contamination of groundwater.

To answer the three points we believed would be true, two out of three isn't bad. In addition, apply high rates of N (2 lbs. N/ 1,000 sq.ft.) did not increase the amount of N leaching when fertilizing in late spring and early fall.

Marty Petrovic, Ph.D., is a Professor of Turfgrass Science, Department of Horticulture, at Cornell University. ■

Rain Bird
encourages
The Intelligent
Use of Water™ every day,
and supports the
Irrigation Association's
Smart Irrigation Month.

The Intelligent Use of Water™

Water. It's what keeps the world alive. As the world's largest manufacturer of irrigation products, we believe it is our responsibility to develop technologies that use water efficiently. Over the past seven decades, our pioneering efforts have resulted in more than 130 patents. From central control systems and automatic shut-off devices to pressure regulating components and low volume drip irrigation, Rain Bird designs products that use water wisely. And our commitment extends beyond products to education, training and services for our industry and our communities.

The need to conserve water has never been greater. We want to do even more, and with your help, we can. Visit www.rainbird.com for more information about The Intelligent Use of Water.™

RAIN BIRD®



Using inorganic soil amendments

By George Toma

In my travels and working with soils over the past 60 years I have seen a number of soil amendment products. In the early days Lusoil was the only soil amendment product available. Later Danny Litwhiler, former player for the Cincinnati Reds, introduced another soil amendment. Danny's product was made from calcined clay. My how the times have changed.

Inorganic soil amendments have been gaining in popularity over the past few years. These products are used in golf course greens, tee boxes, and fairways, sports field rootzone mixes, and topdressing applications. Interest in inorganic rootzone amendments has been amplified due to the potential potable water supply shortages, the increased use of reclaimed water as well as concerns regarding runoff. Until recently, peat was about the only amendment allowed by the USGA for use in

greens. These inorganic products may be better suited to rootzone construction or manipulation because they are less prone to biological degradation and may maintain the original rootzone physical properties longer than organics.

Three types of popular inorganic soil amendments that I am familiar with include calcined clays, zeolite, and diatomaceous earth.

Calcined clays: Marketed as porous ceramics, these are products that have been heat treated or calcined at a very high temperature. The heating increases the structural integrity. Once calcined, they are often screened to a uniform particle size to be used in various rootzones. They do possess a high inherent water-holding capacity that is the result of small internal pores. This stored water may be available to plants. They also have some nutrient holding capacity.

Above: placing an inorganic soil amendment at a NFL practice facility.

INFORMATION PERTAINING TO THE AVERAGE PARTICLE SIZE IN YOUR ROOTZONE MIX MUST BE KNOWN BEFORE ADDING SOIL AMENDMENTS TO AN EXISTING SOIL PROFILE.

Zeolites: Long used in removing environmental pollutants, these have a strong affinity for cations or high CEC, which is good for holding on to nutrients. Be careful in the selection as some zeolites have rather high residual sodium contents that may be harmful to turfgrass.

Diatomaceous Earth (DE): These products are made up of the deposits of diatom shells that have a high degree of internal pore space. DE will retain a significant amount of water. I am very familiar with volcanic/diatomaceous deposits that not only

absorb a considerable amount of water but also make the majority of this stored moisture available to the plant.

There have been numerous studies conducted at universities across the country and these studies should be reviewed before considering what type of amendment should be used. Particle size, water retention and release, and nutrient holding capacity should be taken into consideration when selecting the desired inorganic soil amendment. Also helpful is consultation with an accredited soil-testing laboratory.



George Toma, right, has seen it all when it comes to sports turf amendment products.

WESTERN
Pozzolan



ENGINEERED SOIL PROFILES

westernpozzolan.com

Western Pozzolan produces a wide array of inorganic porous ceramic soil amendments and infield conditioners. Lassenite soil amendments are designed to provide you with the ability to change the characteristics of the soil profile and produce the most desirable balance between air and water filled pore space. This reduces compaction, increasing space available to give the plant moisture, producing deeper, healthier roots.

online seminar
web address:

<http://OneRS.hotims.com/>
9931-305

THE MAJORITY OF PROBLEMS WITH TURF ARE ROOTZONE PROBLEMS THAT CAN BE SOLVED BY DEVELOPING OPTIMUM SOIL CONDITIONS. INORGANIC SOIL AMENDMENTS ARE ANOTHER TOOL THAT CAN BE USED TO PRODUCE POSITIVE RESULTS.

Particle size considerations

Too many coarse or fine particles are undesirable. Particle size significantly affects the amount of porosity and water retention within a rootzone. Fine particles contain less macropores or air filled pores than coarse sands but tend to retain more water than coarse sands. Coarse particles tend to drain more quickly but will exhibit more air filled porosity. Information pertaining to the average particle size in your rootzone mix must be known before adding soil amendments to an existing soil profile.

Water retention

The growth of most plants is closely related to the amount of available moisture. Many inorganic soil amendments are known for their ability to absorb a considerable amount of moisture. The question is: "How much of this stored water is plant available?" If the amendment added to the rootzone releases its stored moisture too quickly, this may lead to droughty conditions or localized dry spots. If the amendment does not release this stored moisture and retain too much water, this will result in poor draining rootzones and cause excessive soil wetness.

Cation exchange capacity

Many inorganic soil amendments have a strong affinity for cations. High cation exchange capacities may allow for a reduction in the use of fertilizers but may also have high residual salt contents. With the increased use of reclaimed water (which may contain elevated sodium levels) on the rise, use of these amendments may prove harmful to turfgrasses.

Desirable criteria for the ideal turfgrass rootzone consists of the following:

- 1) Drainage
- 2) Resistance to compaction
- 3) Moisture retention/release
- 4) Oxygen retention
- 5) Nutrient retention
- 6) Microbial population
- 7) Temperature stability
- 8) Proper porosity balance

Nearly all of the desirable rootzone characteristics are directly tied to the proper balance of porosity.

It's hard to improve on Mother Nature. Mother Nature's perfect rootzone is made up of 50% solids and the remaining 50% consisting of

capillary or water-holding capacity and non-capillary or aeration porosity. Capillary porosity is made up of small pores that hold water against the force of gravity. Depending upon the type of amendment, much of this retained moisture is available for plant use. Non-capillary is made up of larger pores that, when drained, provide for a source of oxygen required for root growth. The number one deficiency in most rootzones is the lack of air. Balancing and manipulating porosity in the soil can be achieved by the application of inorganic amendments.

Application

Incorporation of soil amendments during the construction phase of the soil profile is usually the best way to go. Rootzone characteristics including hydraulic conductivity, water-holding capacity, bulk density, particle size, and porosity can be pre-determined before construction. Inorganic amendments can also be applied during aerification and topdressing or pre-blended with your choice of sands. In addition application via hydraulic injection such as the hydraulic injection equipment of drill and fill may also be used. Several post-aerification applications are required to produce the desired results in terms of water-efficiency and improved growth characteristics.

Many times problems like brown spots or fungus are treated with water or fungicide and the turf manager is really treating symptoms and not causes. The majority of problems with turf are rootzone problems that can be solved by developing optimum soil conditions. Inorganic soil amendments are another tool that can be used to produce positive results.

Much like sports teams and individual players, some products meet the needs of their marketplace and thrive, while others fall short and ultimately disappear. My advice is to thoroughly research the needs of your individual circumstances. Don't be shy; experiment with a number of product options. Evaluate the results of each one. Then make an informed decision. I certainly have my favorites. Here's hoping you find yours and that it meets your needs . . . AND THEN SOME!

George Toma, one of the founders of the Sports Turf Managers Association, is the most famous turf manager in the country, and has worked on the turf for every Super Bowl.

This article was sponsored by Western Pozzolan. ■

Can you identify this sports turf problem?



Problem: Brown parallel lines running across soccer field

Turfgrass Area: High School Athletic Field

Location: Miramar, FL



Grass Variety: 419 Bermudagrass

Answer to

**John Mascaro's Photo
Quiz on Page 41**

*John Mascaro is President
of Turf-Tec International*

The Evolution of Clay.



www.stabilizersolutions.com
1.800.336.2468 local: 602.225.5900



Fill in 123 on reader service form or visit <http://oners.hotims.com/12047-123>

Managing skinned areas

By Bruce Suddeth

When I was asked to write about infield skin management I thought, "Who, me?" How could I write such an article? I don't have the greatest of literary skills, I'm just a landscaper/field manager. Besides, I read every book and article I can find to learn from others how to manage skin areas. The list includes pros like George Toma and Floyd Perry, and that list goes on and on. Our field management team also puts great trust in the South Carolina STMA chapter membership's wealth of knowledge. So how can I write an article about skin management when it is an ongoing learning curve for me?

After much procrastination, I began thinking of what we as "dirt managers" do. Then I saw the light. Here at the University of South Carolina Upstate (USC Upstate) I'm very fortunate to have the situation and resources that we do.

I'd have a safe bet to say that Coach Chris Hawkins puts in more time and is more hands-on with his facility than any coach in the country. You'll see why later.

Composition

Six years ago, our skin area was built of 50% clay, 10% silt, and 40% sand. The field was then amended with three tons of calcined clay conditioner. Since that time more sand and conditioner has been added; the current composition is in the area of 45% clay, 10% silt, and 45% sand. This ratio gives Coach Hawkins the consistency he needs for his team.

Clay bricks are installed in the pitcher's circle and batter's box areas. The bricks are topped off with bagged clay to achieve consistency. The bullpen areas are constructed of the same materials in an attempt to simulate the field's playing surface. Irrigation heads as well as a quick coupler hose for hand watering have been installed in the skin area.

Pre-season prep varies depending on how extensively the skin needs to be repaired. This work typically happens during late November and December. Samples are taken of the skin area and a "cup test" performed to determine the consistency of the soil. We have determined a happy medium between drainage/firming up and moisture retention for playability. This test dictates whether more sand or clay should be added to the skin area.

Whether it is sand or clay the materials are spread with a small top-dresser to achieve consistent results. Using a pulverizer, the materials



Coach Chris Hawkins (left) and Bruce Suddeth.

are incorporated into the existing soil profile. A leveling bar is then used to smooth the soil composition so as to keep the grade (1%) for surface drainage.

Once the field is prepped to Coach Hawkins' satisfaction all play is suspended until practice begins mid-January. During the summer growing season we address any lip area issues. We typically cut two widths 16-inches wide with the sod cutter between second and third bases. We remove the built-up soil and return to the stockpile, then relay the sod. Edges are tamped so as not to have a large transition between skin and turf. In addition, the edges are cut once per week during the growing season. A sharp edge makes the turf and manicured skin stand out.

Tournament, game-day prep

Tournament and game-day prep are for the most part the same. Coach Hawkins applies water to the skin area depending on the nature of the tournament or game. For tournaments the field will be in service for extended hours if not days. Knowing this, Coach Hawkins monitors the amount of water applied to the skin depending on the weather, length of play, and time of year. Coach also takes into account the competition and regulates the skin speed based on the amount of moisture applied. It may

also be necessary to water the skin area between games to maintain a constant playing surface throughout the tournament.

For tournament play the field prep begins the day before. The skin turf interface is blown with a backpack blower to move any conditioner in the turf back into the skin. Water is then applied to the skin to the point of saturation, allowed to stand until absorbed, and the field nail-dragged to loosen the top of the soil layer.

Coach Hawkins also likes to mat (cocoa) drag the skin to break up any clumps and level the area for a more finished look. When dragging the skin he likes to alternate start and stop points at baseline to baseline and inside to outside. This reduces the amount of movement in the skin profile.

It is important to start and stop approximately a foot away from the turf areas to eliminate contamination with the skin soils. These areas should be hand-raked. Another important note is to lift the drag before leaving the skin so as not to pull soil into the turf.

Our practice and game-day prep is much the same as above. We water the morning of practice or game similar to tournament prep. The skin is then dragged with nail and mat for a smooth finish followed up by hand raking along the edges. After and before every event or practice the bullpens are hand raked to keep them looking ready for play.

Due to the amount of effort in keeping Cyrill Softball Stadium ready for play at all times, a constant look at the weather is necessary. During periods of practice, game-day, or tournaments it may be necessary to tarp the field to preserve the skin.

The preparation and work of the skin is not the hard part but the planning, scheduling, coordination, and communication that must happen that is difficult.

Bruce H. Suddeth is director of landscape services for the University of South Carolina Upstate, Spartanburg. He would like to thank USC Upstate Landscape Services, Shurburt CampusScapes, and especially Coach Chris Hawkins for their contributions in winning the 2006 STMA College Softball Field of the Year. ■

www.sportsturfmanager.org



Infield Custom Blends



Vitri-Cal Clay



Diamond Clay Conditioner



Calcined Clay (Surface)



Mound Clay Fortifier



Diamond Sands

We've got a conditioner for every application. Which means less maintenance, less rain-outs, less wind erosion, less complaints and safer conditions at any level of play



MAR-CO
CLAY PRODUCTS INC.
1-800-950-2555
www.marcoclay.com

Fill in 124 on reader service form or visit <http://oners.hotims.com/12047-124>

AerWay®

AerWay® delivers Speed

AerWay® can aerate the typical playing field in less than 30 minutes, without taking it out of play. More frequent aeration treatments reduce compaction, speeds drying of the field when it rains, and makes more efficient use of water when it's dry.



for more information call

1-800-457-8310

Advanced Aeration Systems

www.aerway.com email: aerway@aerway.com

Fill in 125 on reader service form or visit <http://oners.hotims.com/12047-125>



Selecting an **artificial surface**

By John Fik, CSFM

Artificial surfaces have been the “in” thing now for a number of years. High schools are getting state dollars to finance new stadiums and most have been with an artificial surface. I have also seen a large increase of high-end prep schools going to synthetic turf.

I think the first questions I would ask before selecting an artificial surface are “Are you sure you need one?” and “How much

funding do you have secured?” and maybe “What are the factors driving this conversion?” Is it lack of space, lack of technical knowledge by the grounds department to handle a sand base field or a native field, too many sporting activities, concerts, band practices? Is there an increase in summer camps, club, or intramural teams?

If you are trying to play Div 1 and/or Div 3 lacrosse in the northeast like Hobart (men’s) and William Smith (women’s) Colleges in

Geneva, NY are, then it is imperative to have a synthetic field. Our first home game is usually the first or second week of March, which in a normal snow year finds the grass under 2 feet of snow. Most field hockey programs like to play on a synthetic surface; however, before our conversion to synthetic, our women’s team won three national championships on our natural grass field.

The maintenance was very high on that field because we were mowing a k-blue/rye