

Beat insects in 2006

By Daniel A. Potter

ports turf managers need to maintain safe, attractive, playable athletic surfaces regardless of whether their fields serve recreational players or provide the competitive arena for varsity or professional-level athletes. While insects are rarely the first motivation for field management decisions, certain pests can be highly destructive when outbreaks occur. Biting or stinging insects can also be a hazard when they occur on or near playing fields. Use of insecticides in public areas, especially where children are present, is an increasingly volatile issue, requiring field managers to manage pests by means that are non-hazardous to players, bystanders, and the environment.

Fortunately the past 15 years saw the advent of new kinds of insecticides that work selectively against pest insects, are applied at low use rates, and pose low hazard to people and the environment. Many of these products are more versatile and effective than past ones. Let's look at current trends in turf insecticides and what may be on the horizon, and also some non-chemical control options for insect pests of sport fields.

Grubs and billbugs

Root-feeding white grubs are the most destructive insects to sport fields in the cool-season and transitional turfgrass zones. Damage from grubs causes turf to die in irregular patches that can be lifted or rolled back like a loose carpet. Grub damage is worst in late summer and autumn. Skunks and other varmints may dig up the turf to feast on the grubs. Masked chafer and Japanese beetle grubs are abundant throughout most of the eastern and central U.S. Asiatic garden



Turf damaged by white grubs can be rolled back like a carpet. Photo by Daniel Potter.

beetle and Oriental beetle are troublesome in the Northeast, and European chafer grubs occur from the New England states west across New York to northern Ohio and southern Michigan.

Green June beetle grubs are abundant in the transitional climatic zone. They feed more on decaying organic matter than on living roots, but damage turf by burrowing, uprooting the grass, and pushing up mounds of soil. They commonly are associated with bermudagrass fields, sites with high organic matter, or areas where manure or other organic fertilizers have been used.

Registration of Merit (imidacloprid) and MACH 2 (halofenozide) revolutionized grub control during the 1990s, and recent (2005) registration of Arena (clothianidin) provided a third powerful product for preventive grub control. Residues of Merit, MACH 2, and Arena remain

active for as long as 2-3 months, providing flexibility in application timing. All three products are available in granular and spray formulations.

Merit and Arena, which belong to a relatively new class of insecticides called chloronicotinyls, selectively disrupt the insect nervous system. MACH 2 mimics the action of the insect molting hormone, causing a premature lethal molt. All three therefore are target-selective, meaning that they have low inherent toxicity except to insects. They usually provide excellent (>90%) control when applied preventively, before egg hatch. Throughout most of the northern two-thirds of the U.S. the optimal treatment window for preventive grub control is mid-June to mid-July. That timing ensures that fresh residues are in the soil just before and during egg hatch, which for grub pests of sport fields mainly occurs in July and early August. Grub treatments should be watered in as soon as possible to move the residues into the root zone. All three preventive products are, however, relatively forgiving even if rainfall or irrigation is unavoidably delayed for a few days.

Preventive grub insecticides do have limitations. None of them works that well against large grubs, or as "rescue" treatments after damage appears. All three products are highly active against masked chafer and Japanese beetle grubs, the most common and widely distributed grub species. MACH 2 seems to be less effective than Merit or Arena against Asiatic garden beetle, European chafer, and green June beetle grubs, but it may be more active than the others against cutworms and other caterpillars.

The downside of preventive control is that the treatment must be made before the extent of infestation is known. Field managers who reserve preventive treatment for high-risk fields (ones having a history of grubs) or who prefer the "wait-and see" approach may have to spot-treat grub-damaged areas in late summer. Dylox (trichlorfon) is the most effective fast-acting soil insecticide for such curative or "rescue" situations. Water it in immediately (or apply just before a good rain if there is no irrigation) and keep people off the field for 24 hours after application. Insect-parasitic nematodes, specifically *Heterorhabditis bacteriophora*, are an option for non-chemical curative control of white grubs.

Avoid rescue treatment for large green June beetle grubs unless you plan for a cleanup. Grubs of that species die on the surface following such treatments!

Billbugs are small weevils (beetles) that lay eggs in turfgrass stems in spring. Their larvae hollow out the stem and crown, and later migrate to the thatch and soil to feed externally on the crown and roots. With heavy infestations, scattered small patches of dead turf resembling dollar spot disease, may merge into large areas of dead

grass in June to August. With billbugs, tufts of straw-colored dead grass are easily pulled out by hand, the hollowed out stems breaking off at the crown and showing bits of fine sawdust-like frass at their base.

Merit, MACH 2, and Arena also work well against billbugs. They systemically control the young larvae within grass stems, as well as older ones in the soil. Applied at high label rate from mid-to-late May, they will preventively control billbugs with residues persisting long enough to control white grubs later in summer. Alternatively, billbugs can be controlled curatively using MACH 2 or Dylox in early June, or a pyrethroid (see below) can be applied in late April or early May to intercept female billbugs before they lay eggs in the stems. Consult turf specialists at your state university for proper timing in your area.



Mole cricket damage. Photo by Eileen Buss, University of Florida

Surface-feeding pests

Several caterpillar pests including sod webworms, cutworms, and fall armyworms can damage sport fields by chewing the grass blades and stems. Sod webworms and cutworms feed from burrows in the soil and thatch whereas fall armyworms cling to the grass plants while chewing them to the ground. Spot treating with a pyrethroid insecticide easily controls these pests. Options include Talstar (bifenthrin), Tempo (cyfluthrin), DeltaGard (deltamethrin), Scimitar (lambda-cyhalothrin), and Astro (permethrin). Pyrethroids work fast and are applied at very low rates. Remember this when comparing costs because the price per gallon seems high until you factor in cost per application. Pyrethroids also work well against chinch bugs and other sucking pests. They aren't effective against grubs because they bind to thatch and don't reach the root zone. Although pyrethroids have low toxicity to mammals and birds, some formulations are labeled as "Restricted Use" because they're toxic to fish. MACH 2 also will control turf caterpillars.

For turf caterpillars, liquid applications work better than granules because the objective is to leave residues on the foliage and upper thatch. Withhold irrigation and mowing for 24 hours after the application.

Conserve (Spinosad), a reduced-risk insecticide derived from a naturally occurring bacterium, is effective against turf-feeding caterpillars and an excellent choice for sites where use of conventional insecticides might be questioned. *Steinernema carpocapsae*, an insect-parasitic nematode, is another option for caterpillar control.

Mole crickets and fire ants

Mole crickets and fire ants are the most important insect pests of southern sport fields. Mole crickets cause extensive damage by tunneling just under the surface, physically uprooting and stressing the grass, leading to thinning and weed encroachment. They are especially damaging to newly seeded or sprigged fields. Tawny mole crickets also feed extensively on the roots, stems, and leaves. Heavily

infested turf has almost no root system and is easily damaged by sport activities. provides season-long control of both pests. Merit is less expensive and gives good control of mole crickets when applied before or at egg hatch (May to June, depending on latitude). It will simultaneously control white grubs, but has little or no efficacy against fire ants. Arena is in the same ballpark as Merit insofar as mole crickets and grubs, and may also have some activity on fire ants. Some field managers spray as-needed with a pyrethroid (e.g., Talstar) to knock back mole crickets and fire ants. Some recent trials suggest that Allectus, a new combination product containing the active ingredients in Merit and Talstar, may give better mole cricket control than either AI alone. For non-chemical control, try insect-parasitic nematodes, *Steinernema scapterisci* (Nematac S), targeting larger mole cricket nymphs and adults.

If fire ants are the main concern, baits are cheaper (\$10 to 15 per acre, per treatment) and can provide seasonal suppression (e.g., for softball season in spring, football season in fall) with a single



Tawny mole cricket. Photo by Eileen Buss, University of Florida.



Fire ants are a hazard and liability risk on Southern sport fields. Photo by B. Drees, Texas A & M University.

infested turf has almost no root system and is easily damaged by sport activities.

The red imported fire ant occurs throughout the southeastern United States. Like stepping on a biological land mine, disturbing a fire ant colony results in a rapid defensive response by worker ants that swarm up legs or other body parts to inflict numerous bites and stings. Fire ant stings cause an intense burning itch, pustule-like sores, and can be life threatening to persons allergic to the venom. Field managers can become entangled in lawsuits should athletes, spectators, or children at play inadvertently stand or fall on the mounds, which may be inconspicuous when during early stages of colony development. The mounds themselves smother grass and cause damage to mowing equipment.

A lot depends on budgets when managing these pests. TopChoice (fipronil) is the premier mole cricket and fire ant control product being used in the southern states. The cost is relatively high, about \$180 to \$220 per acre for one treatment, but a single broadcast application

application, or sustained suppression with periodic re-treatments. Try the "Texas Two-Step" method (<http://fireant.tamu.edu/>). The first step is to broadcast a bait insecticide over the entire field. The second step is to spot-treat individual mounds with an approved mound drench, granule, bait, or dust insecticide. Baits containing fipronil (Firestar), Abamectin (Affirm), fenoxycarb (Logic, Award), and hydramethylnon (Amdro) are effective. Advion (indoxacarb) is the fastest-acting fire ant bait. It costs a bit more and provides no real residual effect, but is an excellent "rescue treatment" for use on sport fields.

Sound management reduces need for insecticides

Good turf management can reduce chemical inputs and costs. For grubs, use tolerant turf (e.g., turf-type tall fescue), manage thatch, mow at a reasonable height, and follow a balanced fertility regime to promote a deep, extensive root system with good recuperative potential. Consult your extension turf specialist for local recommendations. Night-flying adults of some insect pests (e.g., masked chafers,

European chafers, mole crickets) are attracted to lights, so darkening fields when possible during their flight periods may pay dividends. Irrigated turf attracts adult Japanese beetles, masked chafers, and mole crickets, so cutting back on watering during the flights can discourage their egg laying. Tall fescue and perennial ryegrass cultivars that contain fungal endophytes resist billbugs, sod webworms, and chinch bugs. Overseeding with as little as 40% endophytic perennial rye can reduce populations of those pests, and their damage.

Insecticide labels change so read the label to ensure that a product is labeled for your purposes. For more information, the University of Kentucky Entomology Department website has up-to-date information on white grubs (<http://www.ca.uky.edu/agc/pubs/ent/ent10/ent10.htm>) and grub insecticides (<http://www.uky.edu/Ag/Entomology/entfacts/trees/ef441.htm>). The University of Florida Entomology and Nematology Department (<http://entnemdept.ifas.ufl.edu/>) is excellent on mole crickets, and Texas A&M University has a state-of-the-art website on fire ants (<http://fireant.tamu.edu/>).

Daniel A. Potter is Professor at the University of Kentucky. His book *Destructive Turfgrass Insects: Biology, Diagnosis, and Control* is available from Wiley (www.wiley.com) or Amazon (www.amazon.com). ■



Fire ant mound. Photo by Eileen Buss, University of Florida.

You're Always Ahead of the Game with a COVERMASTER® Raincover...

"Very Good... Highly recommended..."

wrote **Dunham Wright**, Facilities Supervisor, Northeast Louisiana Univ., Monroe, LA

Dunham's comments confirm what we hear from the many groundskeepers who use a COVERMASTER® raincover to keep their fields dry and ready for play.

Call us and we'll gladly tell you more.

The COVERMASTER® Advantage...


- Superior in strength and UV resistance
- Outstanding heat reflective properties
- Light weight - easy to handle
- Widest materials for least number of seams
- Largest choice of weights and colors
- Backed by truly dependable warranties

TARP MACHINE VIDEO!

Call, fax or e-mail for a free video, material samples and a brochure.

COVERMASTER™
COVERMASTER
COVERMASTER

MASTERS IN THE ART OF SPORTS SURFACE COVERS



Covers for football and soccer fields are also readily available.



TARP MACHINE™ lets you roll the cover on and off in minutes.



TARPMATE™ roller comes in 3 lengths with safety end caps.

CALL TOLL FREE
1-800-387-5808

covermaster.com

E-MAIL: info@covermaster.com

MEMBER
SportsTurf
MANAGER ASSOCIATION



COVERMASTER INC., 100 WESTMORE DR. 11-D, REXDALE, ON, M9V 5C3 TEL 416-745-1811 FAX 416-742-6837

Circle 120 on reply card or www.oners.ims.ca/5905-120